

Numerical Analysis & Computer Programming

IFoS (IFS) Previous Year Questions
(PYQ) from 2025 to 2009

Ramana sri Sir

IAS, UPSC, IFS, IFoS, CIVIL
SERVICE MAINS EXAMS
MATHS OPTIONAL STUDY
MATERIALS

2025

1. Find the interval in which the root of the equation $xe^x = 1$ lies between 0 and 1, obtained by using three iterations of bisection method. [8 Marks]
2. Perform the operations (i) $(+42) + (-13)$ and (ii) $(-42) - (-13)$ in binary using signed 2's complement representation for negative numbers in 8-bit system. Give the final answer in decimal. [4+4 Marks]
3. Write down the algorithm for finding the integral $I = \int_a^b f(x) dx$ using Simpson's $\frac{1}{3}$ rule. Hence, find $I = \int_0^1 (4x - 3x^2) dx$ taking 10 intervals. Also, find the relative error. [7+8 Marks]
4. In an examination, the number of candidates who secured marks between certain limits was as follows. Estimate the number of candidates getting marks less than 50. [10 Marks]

Marks	0-19	20-39	40-59	60-89	90-99
No. of candidates	41	62	65	50	17

5. Using Runge-Kutta method of fourth order, find y at $x = 0.2$, given that $\frac{dy}{dx} = 3e^x + 2y$, $y(0) = 0$ and $h = 0.1$, correct to four decimal places. [15 Marks]

2024

6. Obtain the following approximate quadrature formula: [8 Marks]
$$\int_0^3 f(x) dx = \frac{3}{8} [f(0) + 3f(1) + 3f(2) + f(3)].$$
7. Answer the following number-system questions. [4+4 Marks]
(i) Convert $(523.0234375)_{10}$ into an equivalent octal number and then convert it to its binary form.
(ii) If $x = (1D2.2)_{16}$ and $y = (52E.02)_{16}$, then find the value of $x + y$ in decimal system.
8. Write down the algorithm for solving the differential equation $\frac{dy}{dx} = f(x, y)$, $y(x_0) = y_0$ numerically by Euler's method with step length h up to $x = x_n = x_0 + nh$. Solve the following differential equation for $x = 1$ with step length $h = 0.2$ by using Euler's method: [6+9 Marks]
$$\frac{dy}{dx} = x^2 + y, y(0) = 1.$$
9. Show that the iteration formula for the Newton-Raphson method for finding the K^{th} root of a positive real number a is [3+7 Marks]
$$x_{n+1} = \frac{1}{K} [(K-1)x_n + \frac{a}{x_n^{K-1}}], \text{ where } K > 0.$$

Use this formula to find $\sqrt[3]{13}$, correct up to three decimal places.

10. Use Gauss-Jordan elimination method to solve the following system [15 Marks]
of equations, correct up to 2 significant figures:
 $3x_1 + x_2 + x_3 = 7,$
 $2x_1 + x_2 + 5x_3 = 13,$
 $x_1 + 4x_2 + x_3 = 9.4.$

2023

11. A function f defined on $[0, 1]$ is such that $f(0) = 0$, $f(\frac{1}{2}) = -1$ and [8 Marks]
 $f(1) = 0$. Find the quadratic polynomial $p(x)$ which agrees with $f(x)$
for $x = 0, \frac{1}{2}, 1$. If $|\frac{d^3f}{dx^3}| \leq 1$ for $0 \leq x \leq 1$, show that
 $|f(x) - p(x)| \leq \frac{1}{12}$ for $0 \leq x \leq 1$.
12. Draw the logic circuit which realises the Boolean function [8 Marks]
 $L = (A + B)(A + C) + C(A + B \cdot C)$ and simplify it. Draw the
simplified circuit also.
13. Find a root of the equation $\sin x + \cos x = 1$, lying in $(0, 2)$, by [10 Marks]
Regula-Falsi method, correct up to four significant digits.
14. Write down the flow-chart of Runge-Kutta method of 4th order to [15 Marks]
find $y(0.8)$ for $\frac{dy}{dx} = xy$, $y(0) = 2$, taking $h = 0.2$. Also solve the
above IVP to find $y(0.4)$ by Runge-Kutta method (4th order).
15. State the sufficient condition for convergence of the Gauss-Seidel [15 Marks]
iteration method and solve the following system of equations by
using this method, correct up to 3 significant digits:
 $6.7x_1 + 1.1x_2 + 2.2x_3 = 20.5,$
 $2.1x_1 - 15x_2 + 8.4x_3 = 28.8,$
 $3.1x_1 + 9.4x_2 - 15x_3 = 22.9.$

2022

16. Given $f(1) = 4$, $f(2) = 5$, $f(7) = 5$ and $f(8) = 4$. Find the value of $f(6)$ [8 Marks]
and also the value of x for which $f(x)$ is maximum or minimum.
17. Answer the following. [4+4 Marks]
(i) If $x = 0.101010101E0001010$ and $y = 0.100010110E0000110$,
then find $x - y$.
(ii) Draw the map of the Boolean function
 $F = x'yz + xy'z' + xyz + xyz'$. Also simplify the function.
18. Write down an algorithm for Simpson's $\frac{1}{3}$ rule. Hence, compute [6+9 Marks]
 $\int_0^1 x^2(1-x) dx$ correct up to three decimal places with step size
 $h = 0.1$ and compare the result with its exact value.

19. Solve the following system of equations by Gauss-Jordan method: [15 Marks]
 $2x + y - 3z = 11,$
 $4x - 2y + 3z = 8,$
 $-2x + 2y - z = -6.$

20. Find the real root of the equation $e^x - 3x = 0$, by Newton-Raphson method, correct up to four decimal places. [10 Marks]

2021

21. From the following table, estimate the number of students who obtained marks between 40 and 46: [8 Marks]

Marks	30-40	40-50	50-60	60-70	70-80
No. of students	32	43	55	40	30

22. Consider the following integers and their 8-bits binary representations: $13 = 00001101$, $20 = 00010100$. Perform the following bitwise operations and express the results in decimal system: [2+2+2+2 Marks]

- (i) $13 \& 20$ (Bitwise AND)
- (ii) $13 | 20$ (Bitwise OR)
- (iii) $13 \wedge 20$ (Bitwise XOR)
- (iv) ~ 20 (Bitwise Complement)

23. Using Regula-Falsi method, find the fourth root of 28 correct to three decimal places. [15 Marks]

24. Write down the algorithm and flowchart of Runge-Kutta method of fourth order to find the numerical solution at $x = 0.8$ for $\frac{dy}{dx} = \sqrt{2(x + y)}$, $y(0.4) = 0.2$. Assume the step length $h = 0.2$. [7+8 Marks]

25. Solve the following system of equations by Gauss-Seidel method, starting with the initial solution $x_0 = y_0 = z_0 = 0$: [10 Marks]
 $20x + y - 3z = 16,$
 $2x + 20y - z = -19,$
 $3x - 2y + 20z = 25.$