

Complex Analysis

**Previous year Questions
from 2025 to 1992**

2025

1. Expand $f(z) = \frac{1}{(z+1)(z+3)}$ in a Laurent series valid for $1 < |z| < 3$. [10 Marks]
2. Use the method of contour integration to prove that [20 Marks]

$$\int_{-\infty}^{\infty} \frac{x^2 - x + 2}{x^4 + 10x^2 + 9} dx = \frac{5\pi}{12}.$$

3. Evaluate the integral $\int_C \frac{e^z}{z^2(e^z + 1)^3} dz$, where $C : |z| = 2$. [15 Marks]

2024

4. If $w = f(z)$ is an analytic function of z , then show that [10 Marks]

$$\left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) \log |f'(z)| = 0.$$

5. If ϕ and ψ are functions of x and y satisfying Laplace equation, then show that [10 Marks]
 $f(z) = p + iq$, $i = \sqrt{-1}$, is an analytic function, where $p = \frac{\partial \phi}{\partial y} - \frac{\partial \psi}{\partial x}$ and
 $q = \frac{\partial \phi}{\partial x} + \frac{\partial \psi}{\partial y}$.

6. Find a function which is analytic inside and on the circle $C : z = e^{i\theta}$, $0 \leq \theta \leq 2\pi$, and has the value [20 Marks]

$$\frac{(a^2 - 1) \cos \theta + i(a^2 + 1) \sin \theta}{a^4 - 2a^2 \cos 2\theta + 1}$$

on the circumference C , where $a^2 > 1$.

7. Locate the poles and their order of the function $f(z) = \frac{1}{z(\sin \pi z) \left(z + \frac{1}{2} \right)}$. [15 Marks]
Also, find the residue of $f(z)$ at these poles.

2023

8. State the sufficient conditions for a function [10 Marks]
 $f(z) = f(x + iy) = u(x, y) + iv(x, y)$ to be analytic in its domain. Hence, show that $f(z) = \log z$ is analytic in its domain and find $\frac{df}{dz}$.

9. Evaluate $\int_0^{2\pi} \frac{\cos 2\theta}{5 + 4 \cos \theta} d\theta$ using contour integration. [20 Marks]

10. Prove that $u(x, y) = e^x(x \cos y - y \sin y)$ is harmonic. Find its conjugate harmonic function $v(x, y)$ and express the corresponding analytic function $f(z)$ in terms of z . [15 Marks]

11. Classify the singular point $z = 0$ of the function $f(z) = \frac{e^z}{z - \sin z}$ and obtain the principal part of its Laurent series expansion. [15 Marks]

2022

12. If $f(z) = u + iv$ is an analytic function of z , and [10 Marks]

$$u - v = \frac{\cos x + \sin x - e^{-y}}{2 \cos x - e^y - e^{-y}},$$

then find $f(z)$ subject to the condition $f\left(\frac{\pi}{2}\right) = 0$.

13. Expand $f(z) = \frac{1}{(z-1)^2(z-3)}$ in a Laurent series valid for the regions [10 Marks]
 (i) $0 < |z-1| < 2$ and (ii) $0 < |z-3| < 2$.

14. Apply the calculus of residues to evaluate [20 Marks]

$$\int_{-\infty}^{\infty} \frac{\cos x}{(x^2 + a^2)(x^2 + b^2)} dx, \quad a > b > 0.$$

15. Evaluate $\int_C \frac{z+4}{z^2+2z+5} dz$, where C is $|z+1-i| = 2$. [15 Marks]

2021

16. Let $c : [0, 1] \rightarrow \mathbb{C}$ be the curve, where $c(t) = e^{4\pi it}$, $0 \leq t \leq 1$. Evaluate the contour integral [10 Marks]

$$\int_c \frac{dz}{2z^2 - 5z + 2}.$$

17. Find the Laurent series expansion of [20 Marks]

$$f(z) = \frac{z^2 - z + 1}{z(z^2 - 3z + 2)}$$

in the powers of $(z+1)$ in the region $|z+1| > 3$.

18. Let f be an entire function whose Taylor series expansion with centre $z = 0$ has infinitely many terms. Show that $z = 0$ is an essential singularity of $f\left(\frac{1}{z}\right)$. [15 Marks]

19. Using contour integration, evaluate the integral [20 Marks]

$$\int_{-\infty}^{\infty} \frac{\sin x}{x(x^2 + a^2)} dx, \quad a > 0.$$

2020

20. Evaluate the integral $\int_C (z^2 + 3z)dz$ counterclockwise from $(2, 0)$ to $(0, 2)$ along the curve C where C is the circle $|z| = 2$ [10 Marks]
21. Using contour integration, evaluate the integral $\int_0^{2\pi} \frac{1}{3 + 2 \sin \theta} d\theta$ [20 Marks]
22. If $v(r, \theta) = \left(r - \frac{1}{r}\right) \sin \theta, r \neq 0$ then an analytic function $f(z) = u(r, \theta) + iv(r, \theta)$ [15 Marks]

2019

23. Suppose $f(z)$ is Analytical function on a domain D in \mathbb{C} and satisfies the equation. $f(z) = (\operatorname{Re} f(z))^2, z \in D$ Show that $f(z)$ is constant in D [10 Marks]
24. Show that an isolated singular point z_0 of a function $f(z)$ is a pole of order m if and only if $f(z)$ can be written in the form $f(z) = \frac{\phi(z)}{(z - z_0)^m}$ where $\phi(z)$ is analytical and non-zero at z_0 . Moreover

$$\operatorname{Res}_{z=z_0} f(z) = \frac{\phi^{(m-1)}(z_0)}{(m-1)!}, m \geq 1$$
 [15 Marks]
25. Evaluate the integral $\int_C \operatorname{Re}(z^2) dz$ from 0 to $2 + 4i$ along the curve C where C is a parabola $y = x^2$ [10 Marks]
26. Obtain the first three terms of the Laurent series expansion of the function $f(z) = \frac{1}{(e^z - 1)}$ above the point $z = 0$ valid in the region $0 < |z| \leq 2\pi$ [10 Marks]

2018

27. Prove that the function: $u(x, y) = (x-1)^3 - 3xy^2 + 3y^2$ is harmonic and find its harmonic conjugate and the corresponding analytic function $f(z)$ in terms of z . [10 Marks]
28. Find the Laurent's series which represent the function $\frac{1}{(1+z^2)(z+2)}$ when
 (i) $|z| < 1$
 (ii) $1 < |z| < 2$
 (iii) $|z| > 2$ [15 Marks]
29. Show by applying the residue theorem that $\int_0^\infty \frac{dx}{(x^2 + a^2)^2} = \frac{\pi}{4a^3}, a > 0$. [15 Marks]

2017

30. Determine all entire functions $f(z)$ such that 0 is removable singularity of $f\left(\frac{1}{z}\right)$. [10 Marks]
31. Using contour integral method, proves that $\int_0^\infty \frac{x \sin mx}{a^2 + x^2} dx = \frac{\pi}{2} e^{-ma}$. [15 Marks]

32. Let $f = u + iv$ be analytic function on the unit disc $D = \{z \in \mathbb{C} : |z| < 1\}$.
Show that $\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0 = \frac{\partial^2 v}{\partial x^2} + \frac{\partial^2 v}{\partial y^2}$ at all points of D . [15 Marks]
33. For a function $f : \mathbb{C} \rightarrow \mathbb{C}$ and $n \geq 1$, let $f^{(n)}$ denote the n^{th} derivative of f and $f^{(0)} = f$. Let f be an entire function such that for some $n \geq 1$, $f^{(n)}\left(\frac{1}{k}\right) = 0$ for all $k = 1, 2, 3, \dots$. Show that f is a polynomial. [15 Marks]

2016

34. Is $v(x, y) = x^3 - 3xy^2 + 2y$ a harmonic function? Prove your claim, if yes find its conjugate harmonic function and hence obtain the analytic function $u(x, y)$ whose real and imaginary parts are u and v respectively. [10 Marks]
35. Let $\gamma : [0, 1] \rightarrow \mathbb{C}$ be the curve $\gamma(t) = e^{2\pi it}$, $0 \leq t \leq 1$ find giving justification the values of the contour integral $\int_{\gamma} \frac{dz}{4z^2 - 1}$. [15 Marks]
36. Prove that every power series represents an analytic function inside its circle of convergence. [20 Marks]

2015

37. Show that the function $v(x, y) = \ln(x^2 + y^2) + x + y$ is harmonic. Find its conjugate harmonic function $u(x, y)$. Also, find the corresponding analytic function $f(z) = u + iv$ in terms of z . [10 Marks]
38. Find all possible Taylor's and Laurent's series expansions of the function $f(z) = \frac{2z - 3}{z^2 - 3z + 2}$ about the point $z = 0$. [20 Marks]
39. State Cauchy's residue theorem. Using it, evaluate the integral $\int_C \frac{e^z + 1}{z(z+1)(z-i)^2} dz$; $C : |z| = 2$. [15 Marks]

2014

40. Prove that the function $f(z) = u + iv$, where $f(z) = \frac{x^3(1+i) - y^3(1-i)}{x^2 + y^2}$, $z \neq 0$; $f(0) = 0$ satisfies Cauchy-Riemann equations at the origin, but the derivative of f at $z = 0$ does not exist. [10 Marks]
41. Expand in Laurent series the function $f(z) = \frac{1}{z^2(z-1)}$ about $z = 0$ and $z = 1$. [10 Marks]
42. Evaluate the integral $\int_0^{\pi} \frac{d\theta}{\left(1 + \frac{1}{2} \cos \theta\right)^2}$ using residues. [20 Marks]

2013

43. Prove that if $be^{a+1} < 1$ where a and b are positive and real, then the function $z^n e^{-a} - be^z$ has n zeros in the unit circle. [10 Marks]
44. Using Cauchy's residue theorem, evaluate the integral $I = \int_0^\pi \sin^4 \theta \, d\theta$ [15 Marks]

2012

45. Show that the function defined by $f(z) = \begin{cases} \frac{x^3 y^5 (x + iy)}{x^6 + y^{10}}, & z \neq 0 \\ 0, & z = 0 \end{cases}$ is not analytic at the origin though it satisfies Cauchy-Riemann equations at the origin. [12 Marks]
46. Use Cauchy integral formula to evaluate $\int_c \frac{e^{3z}}{(z+1)^4} dz$ where c is the circle $|z| = 2$ [15 Marks]
47. Expand the function $f(z) = \frac{1}{(z+1)(z+3)}$ in Laurent series valid for
- (i) $1 < |z| < 3$
 - (ii) $|z| > 3$
 - (iii) $0 < |z+1| < 2$
 - (iv) $|z| < 1$
- [15 Marks]
48. Evaluate by contour integration $I = \int_0^{2\pi} \frac{d\theta}{1 - 2a \cos \theta + a^2}$ $a^2 < 1$ [15 Marks]

2011

49. If $f(z) = u + iv$ is an analytic function of $z = x + iy$ and $u - v = \frac{e^y - \cos x + \sin x}{\cosh y - \cos x}$, find $f(z)$ subject to the condition, $f\left(\frac{\pi}{2}\right) = \frac{3-i}{2}$ [12 Marks]
50. If the function $f(z)$ is analytic and one valued in $|z - a| < R$, prove that for $0 < r < R$, $f'(a) = \frac{1}{\pi r} \int_0^{2\pi} P(\theta) e^{-i\theta} d\theta$, where $P(\theta)$ is the real part of $f(a + re^{i\theta})$ [15 Marks]
51. Evaluate by Contour integration, $\int_0^1 \frac{dx}{(x^2 - x^3)^{\frac{1}{3}}}$ [15 Marks]
52. Find the Laurent series for the function $f(z) = \frac{1}{1 - z^2}$ with centre $z = 1$ [15 Marks]

2010

53. Show that $u(x, y) = 2x - x^3 + 3xy^2$ is a harmonic function. Find a harmonic conjugate of $u(x, y)$. Hence find the analytic function f for which $u(x, y)$ is the real part. [12 Marks]
54. (i) Evaluate the line integral $\oint_C f(z)dz$ where $f(z) = z^2$, C is the boundary of the triangle with vertices $A(0, 0), B(1, 0), C(1, 2)$ in that order.
- (ii) Find the image of the finite vertical strip $R: x = 5$ to $x = 9$, $-\pi \leq y \leq \pi$ of z -plane under exponential function [15 Marks]
55. Find the Laurent series of the function $f(z) = \exp\left[\frac{\lambda}{2}\left(z - \frac{1}{z}\right)\right]$ as $\sum_{n=-\infty}^{\infty} C_n z^n$ for $0 < |z| < \infty$ where $C_n = \int_0^{2\pi} \cos(n\phi - \lambda \sin \phi) d\phi$, $n = 0, \pm 1, \pm 2, \dots$ with λ a given complex number and taking the unit circle C given by $z = e^{i\phi}$ ($-\pi \leq \phi \leq \pi$) as contour in this region. [15 Marks]

2009

56. Let $f(z) = \frac{a_0 + a_1 z + \dots + a_{n-1} z^{n-1}}{b_0 + b_1 z + \dots + b_n z^n}$, $b_n \neq 0$. Assume that the zeros of the denominator are simple. Show that the sum of the residues of $f(z)$ at its poles is equal to $-\frac{a_n}{b_n}$. [12 Marks]
57. If α, β, γ are real numbers such that $\alpha^2 > \beta^2 + \gamma^2$ show that:
$$\int_0^{2\pi} \frac{d\theta}{\alpha + \beta \cos \theta + \gamma \sin \theta} = \frac{2\pi}{\sqrt{\alpha^2 - \beta^2 - \gamma^2}}$$
 [30 Marks]

2008

58. Find the residue of $\frac{\cot z \coth z}{z^3}$ at $z = 0$ [12 Marks]
59. Evaluate $\int_C \left[\frac{e^{2z}}{z^2(z^2 + 2z + 2)} + \log(z - 6) + \frac{1}{(z - 4)^2} \right] dz$ where C is the circle $|z| = 3$. State the theorems you use in evaluating above integral [15 Marks]

2007

60. Prove that the function f defined by
$$f(z) = \begin{cases} \frac{z^5}{|z|^4} & z \neq 0 \\ 0 & z = 0 \end{cases}$$
 is not differentiable at $z = 0$ [12 Marks]

61. Evaluate (by using residue theorem) $\int_0^{2\pi} \frac{d\theta}{1+8\cos^2 \theta}$ [15 Marks]
62. Show that the transformation $w = z^2$ is conformal at point $z = 1+i$ by finding the images of the lines $y = x$ and $x = 1$ which intersect at $z = 1+i$ [15 Marks]

2006

63. Determine all bilinear transformation which map the half plane $\text{Im}(z) \geq 0$ into the unit circle $|w| \leq 1$ [12 Marks]
64. With the aid of residues, evaluate $\int_0^\pi \frac{\cos 2\theta}{1-2a \cos \theta + a^2} d\theta, -1 < a < 1$ [15 Marks]
65. Prove that all the roots of $z^7 - 5z^3 + 12 = 0$ lie between the circles $|z| = 1$ and $|z| = 2$ [15 Marks]

2005

66. If $f(z) = u + i v$ is an analytic function of the complex variable z and $u - v = e^x(\cos y - \sin y)$, determined $f(z)$ in terms of z . [12 Marks]
67. Expand $f(z) = \frac{1}{(z+1)(z+3)}$ in Laurent's series which is valid for
- (i) $1 < |z| < 3$
 - (ii) $|z| < 3$ and
 - (iii) $|z| < 1$
- [30 Marks]

2004

68. Find the image of the line $y = x$ under the mapping $w = \frac{4}{z^2 + 1}$ and draw the same. Find the points where this transformation ceases to be conformal. [12 Marks]
69. If all zeros of a polynomial $P(z)$ lies in a half plane then show that zeros of the derivatives $P'(z)$ also lie in the same half plane. [15 Marks]
70. Using contour integration evaluate $\int_0^{2\pi} \frac{\cos^2 3\theta}{1-2p \cos 2\theta + p^2} d\theta, 0 < p < 1$ [15 Marks]

2003

71. Determine all the bilinear transformations which transform the unit circle $|z| \leq 1$ into the unit circle $|w| \leq 1$ [12 Marks]
72. Discuss the transformation $W = \left(\frac{z-ic}{z+ic} \right)^2$ (c real) showing that the upper half of the W -plane corresponds to the interior of the semi circle lying to the right of imaginary axis in the z -plane. [15 Marks]

73. Use the method of contour integration to prove that $\int_0^\pi \frac{a d\theta}{a^2 + \sin^2 \theta} = \frac{\pi}{\sqrt{1+a^2}}$ ($a > 0$) [15 Marks]

2002

74. Suppose that f and g are two analytic functions on the set ϕ of all complex numbers with $f\left(\frac{1}{n}\right) = g\left(\frac{1}{n}\right)$ for $n = 1, 2, 3, \dots$. Then show that $f(z) = g(z)$ for each z in ϕ [12 Marks]
75. (i) Show that, when $0 < |z-1| < 2$, that function $f(z) = \frac{z}{(z-1)(z-3)}$ has the Laurent series expansion in powers of $(z-1)$ as $\frac{-1}{2(z-1)} - 3 \sum_{n=0}^{\infty} \frac{(z-1)^n}{2^{n+2}}$ [15 Marks]
76. Establish, by contour integration, $\int_0^\infty \frac{\cos(ax)}{x^2+1} dx = \frac{\pi}{2} e^{-a}$ where $a \geq 0$. [15 Marks]

2001

77. Prove that the Riemann zeta function ζ defined by $\zeta(z) = \sum_{n=1}^{\infty} n^{-z}$ converges for $\operatorname{Re} z > 1$ and converges uniformly for $\operatorname{Re} z \geq 1 + \varepsilon$ where $\varepsilon > 0$ is arbitrary small. [12 Marks]
78. (i) Find the Laurent series for the function $e^{1/z}$ in $0 < z < \infty$. Using this expansion, show that $\frac{1}{\pi} \int_0^\pi \exp(\cos \theta) \cos(\sin \theta - n\theta) d\theta = \frac{1}{n!}$ for $n = 1, 2, 3, \dots$ [15 Marks]
- (ii) Show that $\int_{-\infty}^{\infty} \frac{1}{1+x^4} dx = \frac{\pi}{\sqrt{2}}$ [15 Marks]

2000

79. Show that any four given points of the complex plane can be carried by a bilinear transformation to positions 1, -1, k and $-k$ where the value of k depends on the given points. [12 Marks]
80. Suppose $f(\zeta)$ is continuous on a circle C . Show that $\int_C \frac{f(\zeta) d\zeta}{f(\zeta - x)}$, as z varies inside of C , is differentiable under the integral sign. Find the derivative. Hence or otherwise, derive an integral representation for $f'(z)$ if $f(z)$ is analytic on and inside C . [30 Marks]

1999

81. Examine the nature of the function

$$f(z) = \frac{x^2 y^5 (x + iy)}{x^4 + y^{10}}, z \neq 0,$$

$$f(0) = 0$$

In a region including the origin and hence show that Cauchy-Riemann equations are satisfied at the origin but $f(z)$ is not analytic there. [20 Marks]

82. For the function $f(z) = \frac{-1}{z^3 - 3z + 2}$ find the Laurent series for the domain

(i) $1 < |z| < 2$,

(ii) $|z| > 2$.

Show further that $\oint_C f(z) dz = 0$ where C is any closed contour enclosing that points $z = 1$ and $z = 2$. [20 Marks]

83. Show that the transformation $w = \frac{2z + 3}{z - 4}$ transforms the circle $x^2 + y^2 - 4x = 0$ into the straight line $4u + 3 = 0$, where $w = u + iv$. [20 Marks]

84. Use Residue theorem show that $\int_{-\infty}^{\infty} \frac{x \sin ax}{x^4 + a^4} dx = \frac{\pi}{2} e^{-a} \sin a$, ($a > 0$) [20 Marks]

85. The function $f(z)$ has a double pole at $z = 0$ with residue 2, a simple pole at $z = 1$ with residue 2, is analytic at all other finite points of the plane and is bounded as $|z| \rightarrow \infty$. If $f(2) = 5$ and $f(-1) = 2$ find $f(z)$. [20 Marks]

86. What kind of singularities the following functions have?

(i) $\frac{1}{1 - e^z}$ at $z = 2\pi i$

(ii) $\frac{1}{\sin z - \cos z}$ at $z = \frac{\pi}{4}$

(iii) $\frac{\cot \pi z}{(z - a)^2}$ at $z = a$ and $z = \infty$.

In case (iii) above what happens when a is an integer (including $a = 0$)? [20 Marks]

1998

87. Show that the function

$$f(z) = \frac{x^3(1+i) - y^3(1-i)}{x^2 + y^2}, \quad z \neq 0$$

$$f(0) = 0$$

is continuous and $C - R$ conditions are satisfied at $z = 0$, but $f'(z)$ does not exist at $z = 0$

[20 Marks]

88. Find the Laurent expansion of $\frac{z}{(z+1)(z+2)}$ about the singularity $z = -2$. Specify the region of convergence and the nature of singularity at $z = -2$ [20 Marks]

89. By using the integral representation of $f^n(0)$, prove that $\left(\frac{x^n}{n}\right)^2 = \frac{1}{2\pi i} \oint_C \frac{x^n e^{xz}}{nz^{n+1}} dz$, where C is any closed contour surrounding the origin. Hence show that $\sum_{n=0}^{\infty} \left(\frac{x^n}{n}\right)^2 = \frac{1}{2\pi} \int_0^{2\pi} e^{2x \cos \theta} d\theta$ [20 Marks]

90. Prove that all roots of $z^7 - 5z^3 + 12 = 0$ lie between the circles $|z| = 1$ and $|z| = 2$. [20 Marks]

91. By integrating round a suitable contour show that $\int_0^\infty \frac{x \sin mx}{x^4 + a^4} dx = \frac{\pi}{4b^2} e^{-mb} \sin mb$, where $b = \frac{a}{\sqrt{2}}$ [20 Marks]
92. Using residue theorem, evaluate $\int_0^{2\pi} \frac{d\theta}{3 - 2\cos\theta + \sin\theta}$ [20 Marks]

1997

93. Prove that $u = e^x(x \cos y - y \sin y)$ is harmonic and find the analytic function whose real part is u [20 Marks]
94. Evaluate $\oint_C \frac{dz}{z+2}$ where C is the unit circle. Deduce that $\int_0^{2\pi} \frac{1+2\cos\theta}{5+4\cos\theta} d\theta = 0$ [20 Marks]
95. If $f(z) = \frac{A_1}{z-a} + \frac{A_2}{(z-a)^2} + \dots + \frac{A_n}{(z-a)^n}$ find residue at a for $\frac{f(z)}{z-b}$ where A_1, A_2, \dots, A_n, a and b are constants. What is the residue at infinity? [20 Marks]
96. Find the Laurent series for the function $e^{1/z}$ in $0 < |z| < \infty$. Deduce that $\frac{1}{\pi} \int_0^\pi \exp(\cos\theta) \cdot \cos(\sin\theta - n\theta) d\theta = \frac{1}{n!}, (n = 0, 1, 2, \dots)$ [20 Marks]
97. Integrating e^{-z^2} along a suitable rectangular contour show that $\int_0^\infty e^{-x^2} \cos 2bx dx = \frac{\sqrt{\pi}}{2} e^{-b^2}$ [20 Marks]
98. Find the function $f(z)$ analytic within the unit circle, which takes the values $\frac{a - \cos\theta + i \sin\theta}{a^2 - 2a \cos\theta + 1}$, $0 \leq \theta \leq 2\pi$ on the circle. [20 Marks]

1996

99. Sketchy the ellipse C described in the complex plane by $Z = A \cos \lambda t + iB \sin \lambda t$, $A > B$, where t is real variable and A, B, λ are positive constants. If C is the trajectory of a particle with $z(t)$ as the position vector of the particle at time t , identify with justification
(i) The two positions where the acceleration is maximum, and
(ii) The tow positions were the velocity in minimum [20 Marks]
100. Evaluate $\lim_{z \rightarrow 0} \frac{1 - \cos z}{\sin(z^2)}$ [20 Marks]
101. Show that $z = 0$ is not a branch point for the function $f(z) = \frac{\sin \sqrt{z}}{\sqrt{z}}$. Is it a removable singularity? [20 Marks]
102. Prove that every polynomial equation $a_0 + a_1 z + a_2 z^2 + \dots + a_n z^n = 0$, $a_n \neq 0$, $n \geq 1$ has exactly n roots [20 Marks]
103. By using residue theorem, evaluate $\int_0^\infty \frac{\log_e(x^2 + 1)}{x^2 + 1} dx$ [20 Marks]
104. About the singularity $z = -2$, find the Laurent expansion of $(z-3) \sin\left(\frac{1}{z+2}\right)$. Specify the region of convergence and the nature of singularity at $z = -2$ [20 Marks]

1995

105. Let $u(x, y) = 3x^2y + 2x^2 - y^3 - 2y^2$. Prove that u is a harmonic function. Find a harmonic function v such that $u + iv$ is an analytic function of z . [20 Marks]
106. Find the Taylor series expansion of the function $f(z) = \frac{z}{z^2 + 9}$ around $z = 0$. Find also the radius of convergence of the obtained series. [20 Marks]
107. Let C be the circle $|z| = 2$ described counter clockwise. Evaluate the integral $\int_C \frac{\cosh \pi z}{z(z^2 + 1)} dz$ [20 Marks]
108. Let $a \geq 0$. Evaluate the integral $\int_0^\infty \frac{\cos ax}{x^2 + 1} dx$ with the aid of residues [20 Marks]
109. Let f be analytic in the entire complex plane. Suppose that there exists a constant $A > 0$ such that $|f(z)| \leq A|z|$ for all z . Prove that there exists a complex number a such that $f(z) = az$ for all z [20 Marks]
110. Suppose a power series $\sum_{n=0}^\infty a_n z^n$ convergent at a point $z_0 \neq 0$. Let z_1 be such that $|z_1| < |z_0|$ and $z_1 \neq 0$. Show that the series converges uniformly in the disc $\{z : |z| \leq |z_1|\}$ [20 Marks]

1994

111. Suppose that z is the position vector of a particles moving on the ellipse $C: z = a \cos \omega t + ib \sin \omega t$. Where a, b, ω are positive constants, $a > b$ and t is the time. Determine where
(i) The velocity has the greatest magnitude.
(ii) The acceleration has the least magnitude. [20 Marks]
112. How many zeros does the polynomial $p(z) = z^4 + 2z^3 + 3z + 4$ possess in (i) the first quadrant, (ii) the fourth quadrant [20 Marks]
113. Test of uniform convergence in the region $|z| \leq 1$ the series $\sum_{n=1}^\infty \frac{\cos nz}{n^3}$ [20 Marks]
114. Find Laurent series for
(i) $\frac{e^{2z}}{(z-1)^3}$ about $z = 1$,
(ii) $\frac{1}{z^2(z-3)^2}$ about $z = 3$ [20 Marks]
115. Find the residue of $f(z) = e^z \operatorname{cosec}^2 z$ at all its poles in the finite plane. [20 Marks]
116. By means of contour integration, evaluate $\int_0^\infty \frac{(\log_e u)^2}{u^2 + 1} du$ [20 Marks]

1993

117. In the finite z -plane, show that the function $f(z) = \sec\left(\frac{1}{z}\right)$ has infinitely many isolated singularities in a finite interval which includes 0. [20 Marks]

118. Find the orthogonal trajectories of the family of curves in the xy -plane defined by $e^{-x}(x \sin y - y \cos y) = \alpha$ where α is real function [20 Marks]
119. Prove that (by applying Cauchy Integral formula or otherwise) $\int_0^{2\pi} \cos^{2n} \theta \, d\theta = \frac{1.3.5 \dots (2n-1)}{2.4.6 \dots 2n} 2\pi$ where $n = 1, 2, 3, \dots$ [20 Marks]
120. If c is the curve $y = x^3 - 3x^2 + 4x - 1$ joining the points $(1, 1)$ and $(2, 3)$ find the value of $\int_c (12z^2 - 4iz) \, dz$ [20 Marks]
121. Prove that $\sum_{n=1}^{\infty} \frac{z^n}{n(n+1)}$ converges absolutely for $|z| \leq 1$ [20 Marks]
122. Evaluate $\int_0^{\infty} \frac{dx}{x^6 + 1}$ by choosing an appropriate contour [20 Marks]

1992

123. If $u = e^{-x}(x \sin y - y \cos y)$, find v such that $f(z) = u + iv$ is analytic. Also find $f(z)$ explicitly as function of z [20 Marks]
124. Let $f(z)$ be analytic inside and on the circle C defined by $|z| = R$ and let $z = er^{i\theta}$ be any point inside C . Prove that $f(er^{i\theta}) = \frac{1}{2\pi} \int_0^{2\pi} \frac{(R^2 - r^2)f(Re^{i\phi})}{R^2 - 2Rr \cos(\theta + \phi) + r^2} d\phi$ [20 Marks]
125. Prove that all the roots of $z^7 - 5z^3 + 12 = 0$ lie between the circle $|z| = 1$ and $|z| = 2$. [20 Marks]
126. Find the region of convergence of the series whose n^{th} term is $\frac{(-1)^{n-1} z^{2n-1}}{(2n-1)!}$ [20 Marks]
127. Expand $f(z) = \frac{1}{(z+1)(z+3)}$ in a Laurent series valid for
 (i) $|z| > 3$
 (ii) $1 < |z| < 3$
 (iii) $|z| < 1$ [20 Marks]
128. By integrating along a suitable contour evaluate $\int_0^{\infty} \frac{\cos mx}{x^2 + 1} \, dx$ [20 Marks]